


# Atherogenic indices in clinical practice and biomedical research: A short review

Hussein A. Abid<sup>1</sup>, Zainab S. Abid<sup>2</sup> and Shireen A. Abid<sup>3</sup>

<sup>1</sup>Department of Medical Laboratory Technology, Technical Institute of Baquba, Middle Technical University, Baqubah, 32001, Iraq

<sup>2</sup>Ministry of Health, Health Directorate of Diyala, Baqubah, 32001, Iraq

<sup>3</sup>Department of Chemistry, Faculty of Science, University of Diyala, Baquba, 32001, Iraq



**Received** 20-03-2021

**Revised** 22-04-2021

**Accepted** 10-05-2021

**Published** 23-05-2021

**Corresponding Author**

Hussein A. Abid

[hussainaltameemy@yahoo.com](mailto:hussainaltameemy@yahoo.com)

Department of Medical  
Laboratory Technology, Te

**DOI** <https://doi.org/10.47419/bjbabs.v2i02.52>

**Pages:** 60-70

Distributed under  
the terms of the Creative  
Commons  
Attribution-NonCommercial 4.0  
International (CC-BY-NC 4.0),  
which permits use for any  
non-commercial purpose,  
distribution, and reproduction in  
any medium, provided that the  
original work is properly cited.

**Copyright:** © 2021 Hussein A.  
Abid, Zainab S. Abid, Shireen A.  
Abid

## OPEN ACCESS

## ABSTRACT

Cardiovascular diseases (CVD) represent a major cause of mortality and morbidity worldwide. To date, many physicians still requesting traditional lipid profile tests (TG, TC, HDL-C, and LDL-C) to confirm the clinical diagnosis related to CVD. However, using these tests may be inadequate for predicting CVD risk, especially in intermediate risk. For better clinical practice, laboratory diagnostic alternatives should frequently be evaluated and developed by physicians and laboratory scientists. In this review, we sought to focus on the benefits of lipid ratios (CRI-, CRI-II, AIP, AC, and CHOLindex) in supporting clinical diagnosis and how they can be calculated. A literature search in reputed databases (PubMed and Scopus) was performed to attain this aim, and peer-reviewed research articles were included to conduct this review. Short theoretical and practical notes about each index were accordingly included, along with calculation formulas. Thus, the current article can assist new researchers, and young physicians review what supports their knowledge in managing early CVDs.

**Keywords** atherogenic coefficient, atherogenic Index of plasma, coronary artery disease, dyslipidemia, lipid ratios

## INTRODUCTION

Cardiovascular diseases (CVDs) remain the leading causes of mortality and morbidity worldwide.<sup>1</sup> Most CVDs such as coronary artery disease (CAD), stroke, myocardial infarction (MI), and peripheral vascular disease (PVD) are developed and frequently attributed to a single risk factor called “dyslipidemia”.<sup>2,3</sup> Dyslipidemia can be defined as an elevation in plasma lipids (low-density lipoproteins cholesterol, LDL-C; triglycerides, TGs; or both) or low levels of high-density lipoprotein cholesterol (HDL-C).<sup>4</sup> Genetics, lifestyle, certain disorders (such as thyroid dysfunction and kidney disease), and drugs are major contributors to cause dyslipidemia.<sup>5</sup> In addition to its high prevalence in economically advanced societies, dyslipidemia has been recently reported to find its way in semi-urban populations.<sup>6</sup>

However, early diagnosis of dyslipidemia before the onset of CVD is a helpful cardiovascular preventive measure. In the evaluation of dyslipidemia, lipid profiles (total cholesterol, TC; LDL-C; HDL-C; and TGs) are mostly considered, with emphasis majorly on LDC-C as “bad cholesterol”.<sup>7-9</sup> Apart from their efficiency in diagnosis, using LDL-C alone, or HDL-C alone in predicting CVD risk may be inadequate especially in intermediate-risk when a someone have one or more risk factors (elevated blood lipids, high plasma glucose level, high blood pressure and overweight) that exceed desirable levels or a positive family history.<sup>10-12</sup> Thus, suggesting novel biomarkers or more accurate indices by physicians and biomedical scientists is always needed for best clinical practices. Over the past two decades, many reports showed that in cases when the traditional lipid profiles remain normal, several lipid ratios (atherogenic indices) can be diagnostic alternatives that have been shown in predicting the risk of CVD events and the efficacy of therapy.<sup>13</sup> Among them, atherogenic index of plasma (AIP),<sup>14-17</sup> Castelli’s risk index I (CRI-I) and II (CRI-II),<sup>18</sup> atherogenic coefficient (AC),<sup>9,19,20</sup> and cholesterol index (CHOLindex).<sup>21,22</sup> In this review, the benefits of these indices (ratios) in diagnosing CVD were highlighted and how to calculate each index.


## LIPID RATIOS (ATHEROGENIC INDICES)

### Atherogenic index of plasma

Atherogenic index of plasma (AIP) is an unconventional lipid ratio representing the logarithm of the molar ratio of TGs to HDL-C (Figure 1).<sup>23,24</sup> Accumulated evidence showed that AIP is an important predictive index with a positive correlation with CVD. In 2010, Onat et al. prospectively evaluated 2676 middle-aged Turkey adults with clinical and laboratory investigations during about 8 years’ follow-up and found that AIP is a strong risk factor for CAD in that population.<sup>23</sup> Moreover, Wang et al. (2021) tested the correlation between AIP and SYNTAX (angiographic grading tool to assess the complexity of CAD) score in the Chinese population.<sup>25</sup> They concluded that AIP is an independent risk factor for CAD and may assist in CAD prevention. In addition to many reports which have assessed the AIP in different populations to predict CVD in those populations,<sup>13,16,17,26-28</sup> some researchers suggest a positive correlation with other certain conditions such as acquired premature ejaculation and its severity.<sup>29</sup> Accordingly, evidence supports that AIP is now ready to be considered in clinical practice and for research purposes and further evaluated.

### Castelli’s risk indexes (I & II)


Castelli’s risk indexes (also called cardiac risk indexes) are two lipid ratios, the CRI-I is the ratio of TC to HDL-C (Figure 2), while the CRI-II is the ratio of LDL-C to HDL-C (Figure 3), with notable positive associations with CVD risk.<sup>9,18,30</sup> They were reported by



$$\text{AIP} = \log \left( \frac{\text{TGs}}{\text{HDL-C}} \right)$$


**Figure 1** The mathematical equation to calculate the AIP.

William Castelli, at the end of the past century.<sup>18</sup> Later, many reports assessed and confirmed their positive correlation with CVD<sup>18,24,31–33</sup> and some other conditions such as chronic obstructive pulmonary disease (COPD, negative correlation)<sup>34</sup> and erectile dysfunction.<sup>35</sup> CRI-I has been particularly shown to reflect coronary plaques formation and the thickness of intima-media in the carotid arteries of young adults.<sup>36,37</sup> Therefore, physicians should consider CRI-I and II, in clinical practice to predict or assess atherosclerosis and CVD and for further evaluation.



$$\text{CRI-I} = \frac{\text{TC}}{\text{HDL-C}}$$

**Figure 2** Mathematical equation to calculate CRI-I.



$$\text{CRI-II} = \frac{\text{LDL-C}}{\text{HDL-C}}$$

**Figure 3** Mathematical equation to calculate CRI-II.

## Atherogenic coefficient

Atherogenic coefficient (AC) is the ratio of non-HDL cholesterol to HDL-C (Figure 4).<sup>9,19</sup> It is a diagnostic alternative, which has been used in predicting the risk of developing CV events.<sup>38,39</sup> In this context, it could be a useful tool for CVD specialists to predict a patient's atherogenic status and for further assessment of this index.

$$AC = \frac{TC - HDL-C}{HDL-C}$$

**Figure 4** The mathematical equation to calculate AC.

## Cholesterol index (CHOLindex)

Cholesterol index (CHOLindex) is a simple index that predicts the probability of developing CAD with greater accuracy than the other indices.<sup>21,22</sup> It can be calculated based on LDL-C and HDL-C in individuals with plasma TGs less than 400 mg/dL (Figure 5).<sup>22</sup> In cases when TGs concentration is higher than or equal to 400 mg/dL, the very-low-density lipoprotein cholesterol (VLDL-C) (i.g.  $TGs_{(mg)}/5$ ) should also be included (Figure 6).<sup>22</sup>

$$CHOLindex = LDL-C - HDL-C$$

**Figure 5** Mathematical equation to calculate CHOLindex if the plasma TGs <400 mg/dL.

All atherogenic indices can be easily calculated using simple mathematical formulas, which can be applied automatically within automated analytical devices, with no additional costs are required to be paid for calculating them.

$$\text{CHOLindex (if TGs} \geq 400 \text{ mg/dL)} = -\text{LDL-C} - \text{HDL-C} + \frac{\text{TGs}}{5}$$

**Figure 6** Mathematical equation to calculate CHOLindex if the plasma TGs  $\geq 400$  mg/dL.

## NORMAL RANGES

In view of limited uses in clinical practices, the normal ranges of atherogenic indices cannot easily set. However, each index range in healthy individuals and patients without CVD in previous reports, based on country and sample size, is listed in (Table 1).

Index	Country	Lab. Findings No.	Ranges	Reference	
AIP	Brazil	197	-0.001 $\pm$ 0.30	19	
		95	-0.005 $\pm$ 0.269	34	
	India	100	0.26 $\pm$ 0.14	40	
		699	0.06 $\pm$ 0.002	9	
	Turkey	41	-0.79 $\pm$ 0.15	30	
		51	0.28 $\pm$ 0.26	33	
		48	0.35 $\pm$ 0.35	29	
		32	0.54 $\pm$ 0.22	32	
		32	0.62 $\pm$ 0.64	41	
		19	0.52(0.20–0.59)*	31	
CRI-I	Croatia	95	4.15(3.41–5.04)*	34	
		India	100	3.04 $\pm$ 0.65	42
	Italy	56	4.33(0.14) <sup>‡</sup>	20	
		Mexico	49	Men 4.2(3.56–5.13) <sup>†</sup>	42
	81		Women 3.6(3.02–4.27) <sup>†</sup>	42	
	Nigeria	699	3.76 $\pm$ 0.10	9	
		41	2.18 $\pm$ 0.19	30	
		Turkey	127	4.7 $\pm$ 1.1	22
			48	3.19 $\pm$ 2.42	29
			19	4.27(3.41–5.20)*	31
32			4.4 $\pm$ 0.91	32	
51	3.7(2.4–5.9)*	33			
32	2.26 $\pm$ 1.54	41			
CRI-II	Croatia	95	2.66(2.00–3.53)*	34	
		India	100	1.68 $\pm$ 0.56	42
	Italy	56	2.73(0.11) <sup>‡</sup>	20	

*Continued on next page*

Table 1 continued

	Nigeria	699	2.32±0.09	9
		41	1.12±0.18	30
	Turkey	127	2.9±0.8	22
		51	2.3(1.1–4.5)*	33
		48	1.83±1.54	29
		32	2.9±0.72	32
		32	2.46±0.86	41
		19	2.80(2.18–3.52)*	31
AC	Brazil	197	3.25±1.36	19
	Croatia	95	3.15(2.41–4.04)*	34
	India	100	2.04±0.65	42
	Nigeria	699	2.76±0.10	9
		41	1.18±0.19	30
	Turkey	51	2.7(1.4–4.9)*	33
		48	2.49±2.04	29
		32	3.4±0.91	32
		32	1.27±1.54	41
		19	3.27(2.41–4.20)*	31
CHOLindex (mmol/L)	Nigeria	699	0.78±0.04	9

(\*) = Mean(95% CI), (‡) = Mean(SE), (†) = Median (IQR), all other values are presented as Mean±SD.

## CONCLUSIONS

Unconventional lipid ratios and indices (AIP, CRI-I, CRI-II, AC, and CHOLindex) are positively associated with CVD. Accordingly, these ratios can assist in predicting various CVD events. CHOLindex, based on the available literature, maybe the more accurate index when compared with the others. All atherogenic indices can be easily calculated using simple mathematical formulas, which can be applied automatically within automated analytical devices with no additional costs.

## ABBREVIATIONS

AC, atherogenic coefficient; AIP, atherogenic index of plasma; CAD, coronary artery disease; CHOLindex, cholesterol index; COPD, chronic obstructive pulmonary disease; CRI-I, Castelli's risk index I; CRI-II, Castelli's risk index II; CVD, cardiovascular disease; HDL-C, high-density lipoprotein cholesterol; IQR, interquartile range; LDL-C, low-density lipoprotein cholesterol; MI, myocardial infarction; PVD, peripheral vascular disease; SD, standard deviation; SE, standard error; TC, total cholesterol; TG, triglycerides; VLDL, very-low density lipoprotein cholesterol.

## ACKNOWLEDGEMENTS

The authors would like to thank the reviewers and editors who handled this article's original and revised drafts for their efforts and constructive comments.

## DECLARATIONS

### Authors' contributions

All authors have equally contributed to this paper. They reviewed and approved the final draft submitted to this journal. This article is not associated with data.

### Competing interests

The authors have no conflict of interest.

### Ethical approvals

Not applicable.

### Data availability

Not applicable.

### Funding resources

No fund was received.

## REFERENCES

1. Nielsen TRH. Dyslipidemia and reference values for fasting plasma lipid concentrations in Danish/North-European White children and adolescents. *BMC Pediatrics*. 2017;17:116. Available from: [10.1186/s12887-017-0868-y](https://doi.org/10.1186/s12887-017-0868-y).
2. Rached FH, Chapman MJ, Kontush A. An Overview of the New Frontiers in the Treatment of Atherogenic Dyslipidemias. *Clinical Pharmacology & Therapeutics*. 2014;96(1):57–63. Available from: [10.1038/clpt.2014.85](https://doi.org/10.1038/clpt.2014.85).
3. Perk J. European Guidelines on cardiovascular disease prevention in clinical practice (version 2012): The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by re. *European Heart Journal*. 2012;33(13):1635–1701. Available from: [10.1093/eurheartj/ehs092](https://doi.org/10.1093/eurheartj/ehs092).
4. Sriwijitkamol A, Sangsawang T. Type of dyslipidemia and achievement of the LDL-cholesterol goal in chronic kidney disease patients at the University Hospital. *Vascular Health and Risk Management*. 2015;563:563–563. Available from: [10.2147/vhrm.s92461](https://doi.org/10.2147/vhrm.s92461).
5. Chandra KS, Bansal M, Nair T, Iyengar SS, Gupta R, Manchanda SC, et al. Consensus statement on management of Dyslipidemia in Indian subjects. *Indian Heart Journal*.

- 2014;66(Supplement 3):1–51. Available from: [10.1016/j.ihj.2014.12.001](https://doi.org/10.1016/j.ihj.2014.12.001).
6. Okafor CI. The metabolic syndrome in Africa: Current trends. *Indian Journal of Endocrinology and Metabolism*. 2012;16(1):56–66. Available from: [10.4103/2230-8210.91191](https://doi.org/10.4103/2230-8210.91191).
  7. German CA, Shapiro MD. Assessing Atherosclerotic Cardiovascular Disease Risk with Advanced Lipid Testing: State of the Science. *European Cardiology Review*. 2020;15. Available from: [10.15420/ecr.2019.18](https://doi.org/10.15420/ecr.2019.18).
  8. Nelson RH. Hyperlipidemia as a Risk Factor for Cardiovascular Disease. *Primary Care: Clinics in Office Practice*. 2013;40(1):195–211. Available from: [10.1016/j.pop.2012.11.003](https://doi.org/10.1016/j.pop.2012.11.003).
  9. Olamoyegun M, Oluyombo R, Asaolu S. Evaluation of dyslipidemia, lipid ratios, and atherogenic index as cardiovascular risk factors among semi-urban dwellers in Nigeria. *Annals of African Medicine*. 2016;15(4):194–194. Available from: [10.4103/1596-3519.194280](https://doi.org/10.4103/1596-3519.194280).
  10. Superko HR, King S. Lipid Management to Reduce Cardiovascular Risk: A New Strategy Is Required. *Circulation*. 2008;117(4):560–568. Available from: [10.1161/circulationaha.106.667428](https://doi.org/10.1161/circulationaha.106.667428).
  11. Arad Y, Goodman KJ, Roth M, Newstein D, Guerci AD. Coronary Calcification, Coronary Disease Risk Factors, C-Reactive Protein, and Atherosclerotic Cardiovascular Disease Events. *Journal of the American College of Cardiology*. 2005;46(1):158–165. Available from: [10.1016/j.jacc.2005.02.088](https://doi.org/10.1016/j.jacc.2005.02.088).
  12. Greenland P, Smith SC, Grundy SM. Improving coronary heart disease risk assessment in asymptomatic people: role of traditional risk factors and noninvasive cardiovascular tests. *Circulation*. 2001;104(15):1863–1870. Available from: [10.1161/hc4201.097189](https://doi.org/10.1161/hc4201.097189).
  13. Dobiášová M, Frohlich J, Šedová M, Cheung MC, Brown BG. Cholesterol esterification and atherogenic index of plasma correlate with lipoprotein size and findings on coronary angiography. *Journal of Lipid Research*. 2011;52(3):566–571. Available from: [10.1194/jlr.p011668](https://doi.org/10.1194/jlr.p011668).
  14. Fernández-Macías JC, Ochoa-Martínez AC, Varela-Silva JA, Pérez-Maldonado IN. Atherogenic Index of Plasma: Novel Predictive Biomarker for Cardiovascular Illnesses. *Archives of Medical Research*. 2019;50(5):285–294. Available from: [10.1016/j.arcmed.2019.08.009](https://doi.org/10.1016/j.arcmed.2019.08.009).
  15. Cai G, Shi G, Xue S, Lu W. The atherogenic index of plasma is a strong and independent predictor for coronary artery disease in the Chinese Han population. *Medicine*. 2017;96(37):e8058–e8058. Available from: [10.1097/md.00000000000008058](https://doi.org/10.1097/md.00000000000008058).
  16. Nogay NH. Assessment of the correlation between the atherogenic index of plasma and cardiometabolic risk factors in children and adolescents: might it be superior to the TG/HDL-C ratio? *Journal of Pediatric Endocrinology and Metabolism*. 2017;30(9):947–955. Available from: [10.1515/jpem-2016-0479](https://doi.org/10.1515/jpem-2016-0479).
  17. Aguirre M. Triglycerides/High density lipoprotein cholesterol ratio as a cardiometabolic risk marker in children and adolescents from Mérida city, Venezuela.



- Endocrinol Diabetes Nutr (Engl Ed). 2018;65(2):74–83. Available from: [10.1016/j.endinu.2017.10.010](https://doi.org/10.1016/j.endinu.2017.10.010).
18. Castelli WP, Abbott RD, McNamara PM. Summary estimates of cholesterol used to predict coronary heart disease. *Circulation*. 1983;67(4):730–734. Available from: [10.1161/01.cir.67.4.730](https://doi.org/10.1161/01.cir.67.4.730); <https://dx.doi.org/10.1161/01.cir.67.4.730>.
  19. Nunes SOV. Atherogenic index of plasma and atherogenic coefficient are increased in major depression and bipolar disorder, especially when comorbid with tobacco use disorder. *Journal of Affective Disorders*. 2015;172:55–62. Available from: [10.1016/j.jad.2014.09.038](https://doi.org/10.1016/j.jad.2014.09.038).
  20. Brehm A, Pfeiler G, Pacini G, Vierhapper H, Roden M. Relationship between Serum Lipoprotein Ratios and Insulin Resistance in Obesity. *Clinical Chemistry*. 2004;50(12):2316–2322. Available from: [10.1373/clinchem.2004.037556](https://doi.org/10.1373/clinchem.2004.037556).
  21. Ulusoy RE. LDL cholesterol measurement in terms of CHOLINDEX. *Anadolu Kardiyoloji Dergisi/The Anatolian Journal of Cardiology*. 2013;13(6):612. Available from: [10.5152/akd.2013.208](https://doi.org/10.5152/akd.2013.208).
  22. Akpınar O, Bozkurt A, Acarturk E, Seydaoglu G. A new index (CHOLINDEX) in detecting coronary artery disease risk. *Anadolu Kardiyoloji Dergisi/The Anatolian Journal of Cardiology*. 2013;13(4):315–319. Available from: [10.5152/akd.2013.098](https://doi.org/10.5152/akd.2013.098).
  23. Onat A, Can G, Kaya H, Hergenç G. “Atherogenic index of plasma” (log10 triglyceride/high-density lipoprotein–cholesterol) predicts high blood pressure, diabetes, and vascular events. *Journal of Clinical Lipidology*. 2010;4(2):89–98. Available from: [10.1016/j.jacl.2010.02.005](https://doi.org/10.1016/j.jacl.2010.02.005).
  24. Gómez-Álvarez E, Verdejo J, Ocampo S, Ponte-Negretti CI, Ruíz E, Ríos MM. The CNIC-polypill improves atherogenic dyslipidemia markers in patients at high risk or with cardiovascular disease: Results from a real-world setting in Mexico. *IJC Heart & Vasculature*. 2020;29. Available from: [10.1016/j.ijcha.2020.100545](https://doi.org/10.1016/j.ijcha.2020.100545).
  25. Wang L, Chen F, Xiaoqi C, Yujun C, Zijie L. Atherogenic Index of Plasma Is an Independent Risk Factor for Coronary Artery Disease and a Higher SYNTAX Score. *Angiology*. 2021;72(2):181–186. Available from: [10.1177/0003319720949804](https://doi.org/10.1177/0003319720949804).
  26. MR S, HA C, WG E, M A, AG M, CE LS, et al. Use of the triglyceride/high-density lipoprotein cholesterol ratio to identify cardiometabolic risk: impact of obesity. *Journal of Investigative Medicine*. 2017;65(2):323–327. Available from: [10.1136/jim-2016-000248](https://doi.org/10.1136/jim-2016-000248).
  27. Noumegni SR, Nansseu JR, Bigna JJ, Moor VJA, Assah FK, Dehayem MY, et al. Atherogenic index of plasma and 10-year risk of cardiovascular disease in adult Africans living with HIV infection: A cross-sectional study from Yaoundé, Cameroon. *JRSM Cardiovascular Disease*. 2017;6:1–7. Available from: [10.1177/2048004017740478](https://doi.org/10.1177/2048004017740478);
  28. Sapunar J, Aguilar-Farías N, Navarro J, Araneda G, Chandía-Poblete D, Manríquez V, et al. High prevalence of dyslipidemia and high atherogenic index of plasma in children and adolescents. *Revista Medica de Chile*. 2018;146(10):1112–1122. Available from: [10.4067/S0034-98872018001001112](https://doi.org/10.4067/S0034-98872018001001112).

29. Ertas K, Nas C. Relationship between atherogenic indices and acquired premature ejaculation. *Andrologia*. 2021;53(2). Available from: [10.1111/and.13906](https://doi.org/10.1111/and.13906).
30. Igharo OG. Lipid profile and Atherogenic Indices in Nigerians Occupationally Exposed to e-waste: A Cardiovascular Risk Assessment Study. *Maedica*. 2020;15(2):196–205. Available from: [10.26574/maedica.2020.15.2.196](https://doi.org/10.26574/maedica.2020.15.2.196).
31. Tecer D, Sunar I, Ozdemirel AE, Tural R, Kucuksahin O, Dincel AS, et al. Usefulness of atherogenic indices and Ca-LDL level to predict subclinical atherosclerosis in patients with psoriatic arthritis? *Advances in Rheumatology*. 2019;59(1). Available from: [10.1186/s42358-019-0096-2](https://doi.org/10.1186/s42358-019-0096-2).
32. Koca TT, Tugan CB, Seyithanoglu M, Kocyigit BF. The clinical importance of the plasma atherogenic index, other lipid indexes, and urinary sodium and potassium excretion in patients with stroke. *Eurasian Journal of Medicine*. 2019;51(2):172–176. Available from: [10.5152/eurasianjmed.2019.18350](https://doi.org/10.5152/eurasianjmed.2019.18350).
33. Çakırca G. Lipid profile and atherogenic indices and their association with platelet indices in familial Mediterranean fever. *Turk Kardiyoloji Dernegi Arsivi-Archives of the Turkish Society of Cardiology*. 2018;46:184–190. Available from: [10.5543/tkda.2018.93762](https://doi.org/10.5543/tkda.2018.93762).
34. Markelić I, Hlapčić I, Rogić D, Rako I, Samaržija M, Popović-Grle S, et al. Lipid profile and atherogenic indices in patients with stable chronic obstructive pulmonary disease. *Nutrition, Metabolism and Cardiovascular Diseases*. 2021;31(1):153–161. Available from: [10.1016/j.numecd.2020.07.039](https://doi.org/10.1016/j.numecd.2020.07.039).
35. Culha MG, Canat L, Degirmentepe RB, Albayrak AT, Atalay HA, Merder E, et al. The correlation between atherogenic indexes and erectile dysfunction. *Aging Male*. 2020;23(5):1232–1236. Available from: [10.1080/13685538.2020.1749996](https://doi.org/10.1080/13685538.2020.1749996).
36. Nair D, Carrigan TP, Curtin RJ, Popovic ZB, Kuzmiak S, Schoenhagen P, et al. Association of total cholesterol/ high-density lipoprotein cholesterol ratio with proximal coronary atherosclerosis detected by multislice computed tomography. *Preventive cardiology*. 2009;12(1):19–26. Available from: [10.1111/j.1751-7141.2008.00011.x](https://doi.org/10.1111/j.1751-7141.2008.00011.x).
37. Frontini MG, Srinivasan SR, Xu JH, Tang R, Bond MG, Berenson G. Utility of Non-High-Density Lipoprotein Cholesterol Versus Other Lipoprotein Measures in Detecting Subclinical Atherosclerosis in Young Adults (The Bogalusa Heart Study). *The American Journal of Cardiology*. 2007;100(1):64–68. Available from: [10.1016/j.amjcard.2007.01.071](https://doi.org/10.1016/j.amjcard.2007.01.071).
38. Criqui MH, Golomb BA. Epidemiologic aspects of lipid abnormalities. *The American journal of medicine*. 1998;105(1A):48–57. Available from: [10.1016/s0002-9343\(98\)00212-5](https://doi.org/10.1016/s0002-9343(98)00212-5).
39. Bhardwaj S, Bhattacharjee J, Bhatnagar MK, Tyagi S. Atherogenic Index of Plasma, Castelli Risk Index and Atherogenic Coefficient-New Parameters in Assessing Cardiovascular Risk. *International Journal of Pharmacy and Biological Sciences*. 2013;3(3):359–364.
40. Bora K, Pathak MS, Borah P, Hussain MI, Das D. Association of the Apolipoprotein A-I gene polymorphisms with cardiovascular disease risk factors and atherogenic

- indices in patients from assam, Northeast India. *Balkan Journal of Medical Genetics*. 2017;20(1):59–70. Available from: [10.1515/bjmg-2017-0002](https://doi.org/10.1515/bjmg-2017-0002);
41. Çalışkan Z. Lipid profile atherogenic indeces and their relation of epicardial fat thickness and carotid intima media thickness in Celiac disease. *Northern Clinics of Istanbul*. 2019;6:242–247. Available from: [10.14744/nci.2019.54936](https://doi.org/10.14744/nci.2019.54936).
  42. Banik D, S. Evaluation of Anthropometric Indices and Lipid Parameters to Predict Metabolic Syndrome Among Adults in Mexico. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*. 2021;14:691–701.